




# Learn about your immune system's role in fighting cancer

## Your immune system has a role in fighting cancer.


Learning more about your treatment can help you have better conversations with your healthcare team.



**I am on monotherapy:**  
**OPDIVO® (nivolumab)**



**I am on combination therapy:**  
**OPDIVO + YERVOY® (ipilimumab)**



**OPDIVO and YERVOY can cause your T cells to attack healthy cells contained within organs and tissues in any area of your body and can affect the way they work.** These problems can sometimes become serious or life-threatening and can lead to death, and may happen anytime during treatment or even after treatment has ended. You may have more than one of these problems at the same time. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.

Now, review our educational resources to learn at your own pace.

Visit [www.Immuno-OncologyTeach.com](http://www.Immuno-OncologyTeach.com)

✓ **Check off your progress as you continue.**

Cancer and your immune system

How immunotherapy works

Side effects education

Support team information



Please see additional Important Safety Information for OPDIVO and YERVOY on pages 3-4, and U.S. Full Prescribing Information for [OPDIVO](#) and [YERVOY](#).

# Play an active role in your treatment

It's important to notice any changes in your body after you begin treatment and to report them to your healthcare team right away.

Working with your doctor, use this chart to write down how you've been feeling before you start treatment. The descriptions under each organ of the body can help you put your thoughts into words. After treatment begins, your chart will give you a "baseline" to compare with. It may help you become more aware of changes that could be signs of side effects.

Name of Patient \_\_\_\_\_ Name of Doctor or Nurse \_\_\_\_\_ Date (baseline entry) / \_\_\_\_\_ (follow-up visit)

**Liver:**

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

**Hormone Glands:**

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

**Intestines:**

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

**Lungs:**

- new or worsening cough
- chest pain
- shortness of breath

**Kidneys:**

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

**Other Organs:**

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO or YERVOY. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

**Skin:**

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area

**Severe Infusion Reactions:**

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feel like passing out
- fever
- back or neck pain

**YERVOY may be associated with eye problems** with signs and symptoms that include blurry vision, double vision or other vision problems and eye pain or redness.

If you are taking OPDIVO in combination with another medicine, consult your healthcare professional about the potential side effects of that medicine

## Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).

These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO or YERVOY. Your healthcare provider will monitor you for these complications.

These are not all of the possible side effects of OPDIVO or OPDIVO + YERVOY. This is why you need to report any discomfort or changes in the way you are feeling to your Oncology Care Team.

**i** Please see additional Important Safety Information for OPDIVO (nivolumab) and YERVOY (ipilimumab) on pages 3-4, and U.S. Full Prescribing Information for OPDIVO and YERVOY.

## INDICATIONS

OPDIVO® (nivolumab) is a prescription medicine used to treat people with a type of skin cancer called melanoma that has spread or cannot be removed by surgery (advanced melanoma).

OPDIVO is a prescription medicine used in combination with YERVOY® (ipilimumab) to treat people with a type of skin cancer called melanoma that has spread or cannot be removed by surgery (advanced melanoma).

OPDIVO is a prescription medicine used in combination with YERVOY as a first treatment for adults with a type of advanced stage lung cancer (called non-small cell lung cancer) when your lung cancer has spread to other parts of your body (metastatic) **and** your tumors are positive for PD-L1, but do not have an abnormal EGFR or ALK gene.

OPDIVO is a prescription medicine used in combination with YERVOY and 2 cycles of chemotherapy that contains platinum and another chemotherapy medicine, as a first treatment for adults with a type of advanced stage lung cancer (called non-small cell lung cancer) when your lung cancer has spread or grown, or comes back, **and** your tumor does not have an abnormal EGFR or ALK gene.

OPDIVO is a prescription medicine used to treat people with a type of advanced stage lung cancer (called non-small cell lung cancer) that has spread or grown **and** you have tried chemotherapy that contains platinum, and it did not work or is no longer working. If your tumor has an abnormal EGFR or ALK gene, you should have also tried an FDA-approved therapy for tumors with these abnormal genes, and it did not work or is no longer working.

OPDIVO is a prescription medicine used in combination with YERVOY as a first treatment for adults with a type of cancer that affects the lining of the lungs and chest wall called malignant pleural mesothelioma which cannot be removed by surgery.

OPDIVO is a prescription medicine used in combination with YERVOY to treat people with kidney cancer in certain people when their cancer has spread (advanced renal cell carcinoma) and you have not already had treatment for your advanced RCC.

OPDIVO is a prescription medicine used in combination with cabozantinib to treat people with kidney cancer when your cancer has spread (advanced renal cell carcinoma) and you have not already had treatment for your advanced RCC. Please read the Patient Information that comes with cabozantinib.

OPDIVO is a prescription medicine used to treat people with kidney cancer (renal cell carcinoma) when your cancer has spread or grown after treatment with other cancer medications.

OPDIVO is a prescription medicine used to treat adults with a type of blood cancer called classical Hodgkin lymphoma if your cancer has come back or spread after a type of stem cell transplant that uses your own stem cells (autologous), **and** you used the medicine brentuximab vedotin before or after your stem cell transplant, **or** if you received at least 3 kinds of treatment including an autologous stem cell transplant. OPDIVO was approved based on response rate. There is ongoing evaluation of clinical benefit of OPDIVO for this use.

OPDIVO is a prescription medicine used to treat people with head and neck cancer (squamous cell carcinoma) that has come back or spread **and** you have tried chemotherapy that contains platinum and it did not work or is no longer working.

OPDIVO is a prescription medicine used to treat people with bladder cancer (urothelial carcinoma) that has spread or grown and you have tried chemotherapy that contains platinum, and it did not work or is no longer working. OPDIVO was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO for this use.

## Important Safety Information for OPDIVO and OPDIVO + YERVOY

### What is the most important information I should know about OPDIVO + YERVOY?

OPDIVO and YERVOY are medicines that may treat certain cancers by working with your immune system. OPDIVO and YERVOY can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. These problems may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.

OPDIVO is a prescription medicine used to treat adults and children 12 years of age and older with a type of colon or rectal cancer (colorectal cancer) that has spread to other parts of the body (metastatic), is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), **and** you have tried treatment with a fluoropyrimidine, oxaliplatin, and irinotecan, and it did not work or is no longer working. OPDIVO was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO for this use.

OPDIVO is a prescription medicine used in combination with YERVOY to treat adults and children 12 years of age and older, with a type of colon or rectal cancer (colorectal cancer) that has spread to other parts of the body (metastatic), is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR), **and** you have tried treatment with a fluoropyrimidine, oxaliplatin, and irinotecan, and it did not work or is no longer working. OPDIVO in combination with YERVOY was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO in combination with YERVOY for this use.

OPDIVO is a prescription medicine used to treat people with liver cancer (hepatocellular carcinoma) if you have previously received treatment with sorafenib. OPDIVO was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO for this use.

OPDIVO is a prescription medicine used in combination with YERVOY to treat people with liver cancer (hepatocellular carcinoma) if you have previously received treatment with sorafenib. OPDIVO in combination with YERVOY was approved based on response rate and how long patients' responses lasted. There is ongoing evaluation of clinical benefit of OPDIVO in combination with YERVOY for this use.

OPDIVO is a prescription medicine used to treat people with a type of skin cancer called melanoma to help prevent melanoma from coming back after it and lymph nodes that contain cancer have been removed by surgery.

OPDIVO is a prescription medicine used to treat people with cancer of the tube that connects your throat to your stomach (esophageal cancer) if your esophageal cancer is a type called squamous cell carcinoma, **and** cannot be removed with surgery, **and** has come back or spread to other parts of the body after you have received chemotherapy that contains fluoropyrimidine and platinum.

OPDIVO is a prescription medicine used to treat people with cancer of the stomach (gastric), cancer where the esophagus joins the stomach (gastroesophageal junction), and in people with esophageal adenocarcinoma. OPDIVO may be used in combination with chemotherapy that contains fluoropyrimidine and platinum when your cancer cannot be removed with surgery **or** has spread to other parts of the body.

It is not known if OPDIVO is safe and effective in children younger than 12 years of age with MSI-H or dMMR metastatic colorectal cancer, **or** in children younger than 18 years of age for the treatment of any other cancers.

### Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- **Lung problems:** new or worsening cough; shortness of breath; chest pain
- **Intestinal problems:** diarrhea (loose stools) or more frequent bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; severe stomach-area (abdominal) pain or tenderness
- **Liver problems:** yellowing of your skin or the whites of your eyes; severe nausea or vomiting; pain on the right side of your stomach area (abdomen); dark urine (tea colored); bleeding or bruising more easily than normal

*(continued on next page)*

- **Hormone gland problems:** headaches that will not go away or unusual headaches; eye sensitivity to light; eye problems; rapid heartbeat; increased sweating; extreme tiredness; weight gain or weight loss; feeling more hungry or thirsty than usual; urinating more often than usual; hair loss; feeling cold; constipation; your voice gets deeper; dizziness or fainting; changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- **Kidney problems:** decrease in your amount of urine; blood in your urine; swelling in your ankles; loss of appetite
- **Skin problems:** rash; itching; skin blistering or peeling; painful sores or ulcers in the mouth or nose, throat, or genital area
- **Eye problems:** blurry vision, double vision, or other vision problems; eye pain or redness

**Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO and YERVOY. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include**

- Chest pain; irregular heartbeat; shortness of breath; swelling of ankles
- Confusion; sleepiness; memory problems; changes in mood or behavior; stiff neck; balance problems; tingling or numbness of the arms or legs
- Double vision; blurry vision; sensitivity to light; eye pain; changes in eyesight
- Persistent or severe muscle pain or weakness; muscle cramps
- Low red blood cells; bruising

**Getting medical help right away may help keep these problems from becoming more serious.** Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. Your healthcare team may also need to delay or completely stop your treatment if you have severe side effects.

#### **Possible side effects of OPDIVO + YERVOY**

**OPDIVO and OPDIVO + YERVOY can cause serious side effects, including:**

- **See “What is the most important information I should know about OPDIVO + YERVOY?”**
- **Severe infusion reactions.** Tell your healthcare team right away if you get these symptoms during an infusion of OPDIVO or YERVOY: chills or shaking; itching or rash; flushing; shortness of breath or wheezing; dizziness; feel like passing out; fever; back or neck pain
- **Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO or YERVOY. Your healthcare provider will monitor you for these complications.

**The most common side effects of OPDIVO, when used alone, include:** feeling tired; rash; pain in muscles, bones, and joints; itchy skin; diarrhea; nausea; weakness; cough; vomiting; shortness of breath; constipation; decreased appetite; back pain; upper respiratory tract infection; fever; headache; and stomach-area (abdominal) pain.

**The most common side effects of OPDIVO, when used in combination with YERVOY, include:** feeling tired; diarrhea; rash; itching; nausea; pain in muscles, bones, and joints; fever; cough; decreased appetite; vomiting; stomach-area (abdominal) pain; shortness of breath; upper respiratory tract infection; headache; low thyroid hormone levels (hypothyroidism); decreased weight; and dizziness.

**The most common side effects of YERVOY include:** feeling tired; diarrhea; nausea; itching; rash; vomiting; headache; weight loss; fever; decreased appetite; and difficulty falling or staying asleep.

**The most common side effects of OPDIVO, when used in combination with YERVOY and chemotherapy, include:** feeling tired; pain in muscles, bones, and joints; nausea; diarrhea; rash; decreased appetite; constipation; and itching.

**The most common side effects of OPDIVO, when used in combination with cabozantinib, include:** diarrhea; feeling tired or weak; liver problems; rash, redness, pain, swelling or blisters on the palms of your hands or soles of your feet; mouth sores; rash; high blood pressure; low thyroid hormone levels; pain in muscles, bones, and joints; decreased appetite; nausea; change in the sense of taste; stomach-area (abdominal) pain; cough; upper respiratory tract infection.

**The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:** numbness, pain, tingling, or burning in your hands or feet; nausea; feeling tired; diarrhea; vomiting; decreased appetite; stomach-area (abdominal) pain; constipation; pain in muscles, bones, and joints.

These are not all the possible side effects. For more information, ask your healthcare provider or pharmacist. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

**Before receiving OPDIVO or YERVOY, tell your healthcare provider about all of your medical conditions, including if you:**

- have immune system problems such as Crohn’s disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO and YERVOY can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO or YERVOY passes into your breast milk. Do not breastfeed during treatment with OPDIVO or YERVOY and for 5 months after the last dose of OPDIVO or YERVOY.

#### **Females who are able to become pregnant:**

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO or YERVOY.

- You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of OPDIVO or YERVOY. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with OPDIVO or YERVOY. You or your healthcare provider should contact Bristol Myers Squibb at 1-844-593-7869 as soon as you become aware of a pregnancy.

**Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.**

Please see U.S. Full Prescribing Information and Medication Guide for OPDIVO and YERVOY at [www.OPDIVO.com](http://www.OPDIVO.com).

OPDIVO (10 mg/mL) and YERVOY (5 mg/mL) are injections for intravenous (IV) use.



The **OPDIVO with You** program can help. Call **1-855-OPDIVO-1**



© 2021 Bristol-Myers Squibb Company. All rights reserved. OPDIVO®, YERVOY®, Immuno-Oncology Teach™, and the related logos are trademarks of Bristol-Myers Squibb Company.

7356-US-2100165 02/21

